

**CONFIDENTIAL**



**UNIVERSITI TUN HUSSEIN ONN MALAYSIA**

**FINAL EXAMINATION  
SEMESTER I  
SESSION 2012/2013**

**COURSE NAME : ENGINEERING MATHEMATICS III**  
**COURSE CODE : BSM 2913/BWM 20403/BDA24003**  
**PROGRAMME : 2 BDD/BEE/BFF**  
**EXAMINATION DATE : DECEMBER 2012/JANUARY 2013**  
**DURATION : 3 HOURS**  
**INSTRUCTION : ANSWER ALL QUESTIONS**

**THIS QUESTION PAPER CONSISTS OF FIVE (5) PAGES**

**CONFIDENTIAL**

- Q1** (a) Find the local extremum and saddle point (if exists) for the function

$$f(x, y) = 2x^3 + y^3 + 3x^2 - 3y - 12x - 4.$$

(10 marks)

- (b) Use cylindrical coordinates to find the volume of the solid within the cylinder  $x^2 + y^2 = 9$  and between the planes  $z = 1$  and  $y + z = 2$ .

(10 marks)

- Q2** (a) The displacement of a particle at time  $t$  is given by

$$\mathbf{r}(t) = 2 \cos \frac{\pi}{2} t \mathbf{i} + 2 \sin \frac{\pi}{2} t \mathbf{j} - 2 \mathbf{k}, \quad 0 \leq t \leq 2\pi.$$

Find the velocity  $\mathbf{v}(t)$ , the unit tangent vector  $\mathbf{T}(t)$ , the unit normal vector

$\mathbf{N}(t)$ , binormal vector  $\mathbf{B}(t)$  and curvature  $\kappa$ .

(10 marks)

- (b) Find the centroid of the solid of constant density  $\rho$  bounded by the right circular

cone  $z = \sqrt{x^2 + y^2}$  and plane  $z = 1$ . Given that the mass of the solid is  $\frac{\pi\rho}{3}$ .

(10 marks)

- Q3** (a) Using double integrals, find the surface area of the portion of the sphere

$x^2 + y^2 + z^2 = 9$  that lies above the plane  $z = 1$ .

(10 marks)

- (b) Show that the vector field  $\mathbf{F}(x, y, z) = ye^{-x}\mathbf{i} + (ze^y - e^{-x})\mathbf{j} + e^y\mathbf{k}$  is a conservative vector field and find its potential function  $\phi(x, y, z)$ . Hence evaluate

$\int_C ye^{-x} dx + (ze^y - e^{-x}) dy + e^y dz$ , where  $C$  is the path joining the points  $(0,0,0)$  to

$(3,3,3)$ .

(10 marks)

- Q4** (a) Use Green's theorem to evaluate the integral

$$\oint_C \ln(1+2y)dx - \frac{4xy}{1+2y}dy$$

where  $C$  is the boundary of the region enclosed by  $y = x^2$  and  $x = y$ , and curve  $C$  is oriented counterclockwise.

(10 marks)

- (b) Use Stokes' theorem to evaluate the work done by the force field

$$\mathbf{F}(x, y, z) = -2y \mathbf{i} + 2x \mathbf{j} + z^2 \mathbf{k}$$

along curve  $C$  where  $C$  is the boundary of the surface of sphere  $x^2 + y^2 + z^2 = 10$  that lies above the plane  $z = 1$ .

(10 marks)

- Q5** (a) Evaluate the flux of water  $\mathbf{F}$  across the cone  $z = 9 - \sqrt{x^2 + y^2}$  that lies above the plane  $z = 0$ , oriented by an upward unit normal vector, where the velocity vector field  $\mathbf{F}(x, y, z) = 2x \mathbf{i} + 2y \mathbf{j} + 4 \mathbf{k}$  is measured in m/sec.

[Water has the density  $\rho = 1$  unit]

(10 marks)

- (b) Use the Gauss's Theorem to evaluate the flux of vector field  $\mathbf{F}$  across surface  $\sigma$ , where  $\mathbf{F}(x, y, z) = 2x^3 \mathbf{i} + 2y^3 \mathbf{j} + 2z^3 \mathbf{k}$  and  $\sigma$  is the surface of the solid enclosed by upper hemisphere  $z = \sqrt{4 - x^2 - y^2}$  and the plane  $z = 0$ , oriented outward.

(10 marks)

**FINAL EXAMINATION**

SEMESTER / SESSION: SEM II / 2010/2011

COURSE : BDD/BEE/BFF

SUBJECT : ENGINEERING MATHEMATICS III

CODE : BSM 2913/BWM 20403

**Formulae**

**Polar coordinate:**  $x = r \cos \theta$ ,  $y = r \sin \theta$ ,  $\theta = \tan^{-1}(y/x)$ , and  $\iint_R f(x, y) dA = \iint_R f(r, \theta) r dr d\theta$

**Cylindrical coordinate:**  $x = r \cos \theta$ ,  $y = r \sin \theta$ ,  $z = z$ ,  $\iiint_G f(x, y, z) dV = \iiint_G f(r, \theta, z) r dz dr d\theta$

**Spherical coordinate:**  $x = \rho \sin \phi \cos \theta$ ,  $y = \rho \sin \phi \sin \theta$ ,  $z = \rho \cos \phi$ ,  $x^2 + y^2 + z^2 = \rho^2$ ,  
 $0 \leq \theta \leq 2\pi$ ,  $0 \leq \phi \leq \pi$ , and

$$\iiint_G f(x, y, z) dV = \iiint_G f(\rho, \phi, \theta) \rho^2 \sin \phi d\rho d\phi d\theta$$

**Directional derivative:**  $D_{\mathbf{u}} f(x, y) = (f_x \mathbf{i} + f_y \mathbf{j}) \cdot \mathbf{u}$

Let  $\mathbf{F}(x, y, z) = M \mathbf{i} + N \mathbf{j} + P \mathbf{k}$  is vector field, then

$$\text{the divergence of } \mathbf{F} = \nabla \cdot \mathbf{F} = \frac{\partial M}{\partial x} + \frac{\partial N}{\partial y} + \frac{\partial P}{\partial z}$$

$$\text{the curl of } \mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ M & N & P \end{vmatrix} = \left( \frac{\partial P}{\partial y} - \frac{\partial N}{\partial z} \right) \mathbf{i} - \left( \frac{\partial P}{\partial x} - \frac{\partial M}{\partial z} \right) \mathbf{j} + \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \mathbf{k}$$

Let  $C$  is a smooth curve given by  $\mathbf{r}(t) = x(t) \mathbf{i} + y(t) \mathbf{j} + z(t) \mathbf{k}$ ,  $t$  is parameter, then

$$\text{the unit tangent vector: } \mathbf{T}(t) = \frac{\mathbf{r}'(t)}{\|\mathbf{r}'(t)\|}$$

$$\text{the unit normal vector: } \mathbf{N}(t) = \frac{\mathbf{T}'(t)}{\|\mathbf{T}'(t)\|}$$

$$\text{the binormal vector: } \mathbf{B}(t) = \mathbf{T}(t) \times \mathbf{N}(t)$$

$$\text{the curvature: } \kappa = \frac{\|\mathbf{T}'(t)\|}{\|\mathbf{r}'(t)\|} = \frac{\|\mathbf{r}'(t) \times \mathbf{r}''(t)\|}{\|\mathbf{r}'(t)\|^3}$$

$$\text{the radius of curvature: } \rho = 1/\kappa$$

$$\text{Green Theorem: } \oint_C M dx + N dy = \iint_R \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dA$$

$$\text{Gauss Theorem: } \iint_S \mathbf{F} \cdot \mathbf{n} dS = \iiint_G \nabla \cdot \mathbf{F} dV$$

$$\text{Stokes' Theorem: } \oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS$$

**Arc length**

$$\text{If } \mathbf{r}(t) = x(t) \mathbf{i} + y(t) \mathbf{j}, t \in [a, b], \text{ then the arc length } s = \int_a^b \|\mathbf{r}'(t)\| dt = \int_a^b \sqrt{[x'(t)]^2 + [y'(t)]^2} dt$$

$$\text{If } \mathbf{r}(t) = x(t) \mathbf{i} + y(t) \mathbf{j} + z(t) \mathbf{k}, t \in [a, b], \text{ then the arc length } s = \int_a^b \sqrt{[x'(t)]^2 + [y'(t)]^2 + [z'(t)]^2} dt$$

**FINAL EXAMINATION**

SEMESTER / SESSION: SEM II / 2010/2011

COURSE : BDD/BEE/BFF

SUBJECT : ENGINEERING MATHEMATICS III

CODE : BSM 2913/BWM 20403

**Tangent Plane**

$$z - z_0 = f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0)$$

**Extreme of two variable functions**

$$G(x, y) = f_{xx}(x, y)f_{yy}(x, y) - (f_{xy}(x, y))^2$$

Case1: If  $G(a, b) > 0$  and  $f_{xx}(x, y) < 0$  then  $f$  has local maximum at  $(a, b)$ Case2: If  $G(a, b) > 0$  and  $f_{xx}(x, y) > 0$  then  $f$  has local minimum at  $(a, b)$ Case3: If  $G(a, b) < 0$  then  $f$  has a saddle point at  $(a, b)$ Case4: If  $G(a, b) = 0$  then no conclusion can be made.**In 2-D: Lamina**

**Mass:**  $m = \iint_R \delta(x, y) dA$ , where  $\delta(x, y)$  is a density of lamina.

**Moment of mass:** (i) about  $y$ -axis,  $M_y = \iint_R x\delta(x, y) dA$ , (ii) about  $x$ -axis,  $M_x = \iint_R y\delta(x, y) dA$

**Centre of mass,**  $(\bar{x}, \bar{y}) = \left( \frac{M_y}{m}, \frac{M_x}{m} \right)$

**Moment inertia:** (i)  $I_y = \iint_R x^2 \delta(x, y) dA$ , (ii)  $I_x = \iint_R y^2 \delta(x, y) dA$ , (iii)  $I_o = \iint_R (x^2 + y^2) \delta(x, y) dA$

**In 3-D: Solid**

**Mass,**  $m = \iiint_G \delta(x, y, z) dV$ . If  $\delta(x, y, z) = c$ ,  $c$  is a constant, then  $m = \iiint_G dA$  is volume.

**Moment of mass**

(i) about  $yz$ -plane,  $M_{yz} = \iiint_G x\delta(x, y, z) dV$

(ii) about  $xz$ -plane,  $M_{xz} = \iiint_G y\delta(x, y, z) dV$

(iii) about  $xy$ -pane,  $M_{xy} = \iiint_G z\delta(x, y, z) dV$

**Centre of gravity,**  $(\bar{x}, \bar{y}, \bar{z}) = \left( \frac{M_{yz}}{m}, \frac{M_{xz}}{m}, \frac{M_{xy}}{m} \right)$

**Moment inertia**

(i) about  $x$ -axis:  $I_x = \iiint_G (y^2 + z^2) \delta(x, y, z) dV$

(ii) about  $y$ -axis:  $I_y = \iiint_G (x^2 + z^2) \delta(x, y, z) dV$

(iii) about  $z$ -axis:  $I_z = \iiint_G (x^2 + y^2) \delta(x, y, z) dV$